DNA Degradase™

Cat. Nos. E2016 (500 Units)

E2017 (2,000 Units)

Storage: -20 °C




Product Information

Highlights:

- Quick and simple procedure for generating single <u>nucleotides</u> from DNA for quantitative analysis (e.g., wholegenome methylation analysis by HPLC, TLC, etc.)
- 1 hour, single-enzyme digest vs. conventional 6-16 hour multi-step enzyme digestion protocols.

Description:

DNA Degradase $^{\text{TM}}$ from Zymo Research is a nuclease mix that *quickly* and *efficiently* degrades DNA to its individual *nucleotide* components. DNA Degradase $^{\text{TM}}$ is ideal for whole-genome DNA methylation analysis by a number of downstream applications (i.e., HPLC, TLC, etc.). Digestion with the enzyme is performed via a *one-step* procedure that is faster and simpler than other available methods.




Quantitation Thin layer chromatography

Applications HPLC

LC/Mass Spectrometry

DNA Degradase™ degrades DNA to individual <u>nucleotides</u>. Quantitation of nucleotides to detect DNA methylation levels can be assessed by various applications (e.g.- TLC, HPLC, etc.). For detection of DNA methylation via LC/MS, DNA needs to be degraded down to individual <u>nucleosides</u>, see DNA Degradase Plus™ (cat nos. E2020, E2021).



DNA Degradase[™] Efficiently Degrades DNA: 1 μg, 5 μg, and 10 μg of λ DNA incubated with 1 μl (10 U) of DNA Degradase[™] in 25 μl reaction volume (see *Standard Reaction Setup*) and incubated at 37°C for 1 hour.

Product Contents:

| | Cat. No. E2016 | Cat. No. E2017 | Storage |
|--|-------------------|-------------------|---------|
| DNA Degradase™ | 500 units | 2,000 units | -20°C |
| 10X DNA Degradase™ Reaction Buffer (E2016-2) | 1 ml | 1 ml | -20°C |

Storage: Store reagents at -20°C for up to 12 months. Avoid repeated freeze/thawing. Prolonged storage should be ≤ -70 °C.

Enzyme Concentration: 10 units/µl

Unit Definition: One unit (1 U) is defined as the amount of enzyme required to degrade 1 μg of λ DNA in a total reaction volume of 25 μl for 1 hour at 37°C.

Reaction Conditions: DNA Degradase™ in 1X DNA Degradase™ Reaction Buffer. Incubate reaction mixtures at 37°C for ≥1hour.

Inactivation: Heat-inactivate at 70°C for 20 minutes.

Standard Reaction Setup: The setup (below) is an example of a typical DNA DegradaseTM reaction in a 25 μ I final volume. However, the volumes can be adjusted accordingly depending on experimental requirements.

2 μl DNA at 500 ng/μl

2.5 µl 10X DNA Degradase™ Reaction Buffer

1 μl DNA Degradase™ (10 units/μl)

19.5 µl ddH₂O

25 µl Total Volume

Incubate at 37°C for 2 hours (see Note 1, below).

Notes:

1. Complete digestion of DNA.

For *complete* digestion of DNA (gDNA, plasmid, amplified DNA, etc.), reactions should be carried out at 37 °C for \geq 2 hours. Even though 1 unit of DNA DegradaseTM is sufficient to digest 1 µg of DNA at 37 °C in 1 hour, we recommend "overdigesting" with 5-10 fold enzyme to ensure complete digestion of DNA.

2. Sample DNA purity

The digestion reaction can be sensitive to extraneous components from crude DNA preparations or DNA samples carried over from prior reactions. It is recommended that impure preparations of DNA be "cleaned" prior to manipulation (e.g., DNA Clean and Concentrator from Zymo Research).

DNA Degradase™

Cat. Nos. E2016 (500 Units)

E2017 (2,000 Units)

Storage: -20 °C



Product Information

Trademarks and Disclaimers:

™ Trademarks of Zymo Research Corporation.

This product is for research use only and should only be used by trained professionals. It is not intended for use in diagnostic procedures. Wear protective gloves and eye protection. Follow the safety guidelines and rules enacted by your research institution or facility.

Integrity of kit components is guaranteed for up to one year from date of purchase. Reagents are routinely tested on a lot-to-lot basis to ensure they provide maximal performance and reliability.

For Technical Assistance, please contact 1-888-882-9682 or E-mail tech@zymoresearch.com. Satisfaction of all Zymo Research products is guaranteed. If you are not satisfied with this product please call 1-888-882-9682.

Version 1.0.1

| Related Epigenetics Products: | 1 | |
|---|------------------------------|----------------|
| Product Name | Size | Cat. No. |
| BISULFITE TREATMENT OF | | |
| | 50 rxns. | D5001 |
| EZ DNA Methylation™ Kit | 200 rxns. | D5002 |
| | 2 x 96 rxns. 2 x 96 rxns. | D5003 D5004 |
| | 50 rxns. | D5004 |
| EZ DNA Methylation-Gold™ Kit | 200 rxns. | D5006 |
| | 2 x 96 rxns. | D5007 |
| | 2 x 96 rxns. | D5008 |
| EZ DNA Methylation-Direct™ Kit | 50 rxns. | D5020 |
| | 200 rxns. 2 x 96 rxns. | D5021 |
| | 2 x 96 rxns. | D5022 D5023 |
| EZ DNA Methylation-Startup™ Kit | 50 rxns. | D5023 |
| LZ DNA Methylation-Startup - Nit | | D5024 |
| | 50 preps. 200 preps. | D5025 |
| EZ Bisulfite DNA Clean-up Kit™ | 2 x 96 preps. | D5027 |
| | 2 x 96 preps. | D5028 |
| METHYLATED/NON-METHYLATED DI | NA STANDARDS | |
| Jniversal Methylated DNA Standard | 1 set | D5010 |
| <u> </u> | | |
| Universal Methylated Human DNA Standard | 1 set | D5011 |
| Universal Methylated Mouse DNA Standard | 1 set | D5012 |
| Human Methylated and Non-methylated DNA Set | 1 set | D5014 |
| Human HCT116 DKO Non-methylated DNA Standard | 5 µg | D5014-1 |
| Human HCT116 DKO Methylated DNA Standard | 5 μg | D5014-2 |
| Bisulfite Converted Universal Methylated Human DNA Standard | 1 set | D5015 |
| E. coli Non-methylated Genomic DNA | 5 μg | D5016 |
| AMPLIFICATION OF BISULFITE CO | NVERTED DNA | |
| Z To a TM DNIA Dolumono | 50 rxns. | E2001 |
| Zymo <i>Taq</i> ™ DNA Polymerase | 200 rxns. | E2002 |
| Zymo <i>Taq</i> ™ PreMix (2X concentrated) | 50 rxns. | E2003 |
| zymoray ···· Freiviix (2A concentrated) | 200 rxns. | E2004 |
| ANTIBODIES & IMMUNOPRECI | PITATION | |
| Methylated-DNA IP Kit | 10 preps. | D5101 |
| | 50 preps. | D5201 |
| ChIP DNA Clean & Concentrator™ | 50 preps. | D5205 |
| Anti-5-Methylcytosine Monoclonal | 50 μg | A3001-50 |
| Antibody (clone 10G4) | 200 μg | A3001-200 |
| METHYLTRANSFERASI | ES . | |
| | 200 U | E2010 |
| CpG Methylase (M.Sssl) | 400 U | E2011 |
| GpC Methylase (M.CviPI) | 200 U | E2014 |
| SPC Methylase (M.CVIFI) | 1000 U | E2015 |
| DNA FRAGMENTATIO | N | |
| DNA Degradase™ | 500 U | E2016 |
| - | 2000 U 250 U | E2017 |
| DNA Degradase Plus™ | 1000 U | E2020 E2021 |
| | 50 U | E2018-50 |
| DNA Shearase™ | 200 U | E2018-200 |
| OINA OIIEdidae | 50 U & DCC™ | E2019-50 |
| | 200 U & DCC™ | E2019-200 |
| NUCLEOSOME MAPPIN | IG | |
| EZ Nucleosomal DNA Prep Kit | 20 preps | D5220 |
| 5-HYDROXYMETHYLCYTO | SINE | |
| 5-Hydroxymethylcytosine DNA | 5 µg | D5400 |
| 3-i iyuloxyilletiiyicytosiile DivA | | |
| 5-Methylcytosine & 5-Hydroxymethylcytosine DNA Standard Set | 1 set | D5405 |